

Numerical Methods

(MTH4002)

Lecture 06: Numerical Integration


Dr. Kundan Kumar
Associate Professor
Department of ECE



Faculty of Engineering (ITER)
S'O'A Deemed to be University, Bhubaneswar, India-751030
© 2020 Kundan Kumar, All Rights Reserved

Introduction

- Integration is frequently encountered when solving problems and calculating quantities in engineering and science.
- One of the **simplest examples** for the application of integration is the **calculation of the length of a curve**.



- When a curve in the x - y plane is given by the equation $y = f(x)$, the length L of the curve between the points $x = a$ and $x = b$ is given by:


$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

Background

- The general form of a **definite integral** (also called an **antiderivative**) is:

$$I(f) = \int_a^b f(x)dx$$

where $f(x)$, called the **integrand**, is a function of the independent variable x , and a and b are the **limits of the integration** (**definite integration**).




- The value of the integral $I(f)$ is a number when a and b are numbers.
- Graphically, the **value of the integral** corresponds to the shaded **area under the curve** of $f(x)$ between a and b .

Need for numerical integration

- The integrand can be an analytical function or a set of discrete points (tabulated data).
- When the integrand is a mathematical expression for which the antiderivative can be found easily, the value of the definite integral can be determined analytically.
- Numerical integration is needed when analytical integration is difficult or not possible, and when the integrand is given as a set of discrete points.

Numerical Integration Approach

- If the integrand $f(x)$ is an analytical function, the numerical integration is done by using a finite number of points at which the integrand is evaluated.



- One strategy is to use only the end points of the interval, $(a, f(a))$ and $(b, f(b))$.
- This, however, might not give an accurate enough result, especially if the interval is wide and/or the integrand varies significantly within the interval.
- Higher accuracy can be achieved by using a **composite method** where the **interval $[a, b]$** is **divided into smaller subintervals**.



Numerical Integration Approach

- The integral over each subinterval is calculated, and the **results are added** together to give the value of the **whole integral**.
- In all cases, the **numerical integration** is carried out by using **a set of discrete points for the integrand**.
- When the **integrand is an analytical function**, the location of the **points within the interval $[a, b]$** can be defined by the user or is defined by the integration method.
- When the integrand is a given **set of tabulated points** (like data measured in an experiment), the location of the points is **fixed and cannot be changed**.

Numerical Integration Approach

- Various methods have been developed for carrying out numerical integration.
- In each of these methods, a formula is derived for calculating an approximate value of the integral from discrete values of the integrand.
- The methods can be divided into two groups
 - open methods and
 - closed methods.

Numerical Integration Approach




- In numerical methods, a formula is derived for calculating an approximate value of the integral from discrete values of the integrand.
- In closed integration methods, the endpoints of the interval (and the integrand) are used in the formula that estimates the value of the integral.
 - Trapezoidal method
 - Simpson's method
- In open integration methods do not include the endpoints in the formula.
 - Midpoint method
 - Gauss quadrature

An another approach for integration

- There are various methods for calculating the value of an integral from the set of discrete points of the integrand. Most commonly, it is done by using **Newton-Cotes integration formulas**.
- When the original integrand is an analytical function, the Newton-Cotes formula replaces it with a simpler function.
- When the original integrand is in the form of data points, the Newton-Cotes formula interpolates the integrand between the given points.
- Most commonly, as with the trapezoidal method and Simpson's methods, the Newton-Cotes integration formulas are polynomials of different degrees.

An another approach for integration

- The approach for integration is to curve-fit the points with a function $F(x)$ that best fits the points (the function $f(x)$ is must be specified as discrete points).
- In other words, $f(x) \approx F(x)$, where $F(x)$ is a polynomial or a simple function whose antiderivative can be found easily. Then, the integral is evaluated by direct analytical methods from calculus.



$$I(f) = \int_a^b f(x)dx \approx \int_a^b F(x)dx$$

- This procedure requires numerical curve fitting methods for finding $F(x)$, but may not require a numerical method to evaluate the integral if $F(x)$ is an integrable function.

Rectangle and midpoint methods

■ Rectangle method





$$I(f) = \int_a^b f(x) dx \approx f(a)(b-a)$$

$$\text{or } I(f) = \int_a^b f(x) dx \approx f(b)(b-a)$$

■ Composite rectangle method

$$I(f) = \int_a^b f(x) dx \approx h \sum_{i=1}^N f(x_i)$$


Exact integral

Approximating the integral assuming $f(x)=f(a)$ Approximating the integral assuming $f(x)=f(b)$ 

Rectangle and midpoint methods


- Midpoint method

$$I(f) = \int_a^b f(x) dx \approx \int_a^b f\left(\frac{a+b}{2}\right) dx = f\left(\frac{a+b}{2}\right) (b-a)$$



- Composite midpoint method


$$I(f) = \int_a^b f(x) dx \approx h \sum_{i=1}^N f\left(\frac{x_i + x_{i+1}}{2}\right)$$



Trapezoidal method

- A refinement over the simple rectangle and midpoint methods is to use a **linear function to approximate the integrand** over the interval of integration.
- Newton's form of interpolating polynomials with two points $x = a$ and $x = b$, yields:

$$f(x) \approx f(a) + (x - a)f[a, b] = f(a) + (x - a) \frac{[f(b) - f(a)]}{b - a}$$




so we can write

$$\begin{aligned} I(f) &\approx \int_a^b \left(f(a) + (x - a) \frac{[f(b) - f(a)]}{b - a} \right) dx \\ &= f(a)(b - a) + \frac{1}{2}[f(b) - f(a)](b - a) \\ &= \frac{[f(a) + f(b)]}{2}(b - a) \end{aligned}$$

Trapezoidal method

- Simplifying the result gives an approximate formula popularly known as the **trapezoidal rule** or **trapezoidal method**.




- As with the rectangle and midpoint methods, the trapezoidal method can be easily extended to yield any **desired level of accuracy by subdividing the interval $[a, b]$** into subintervals.

$$I(f) \approx \frac{[f(a) + f(b)]}{2} (b - a)$$

Composite Trapezoidal

- The integral over the interval $[a, b]$ can be evaluated more accurately by dividing the interval into subintervals, **evaluating the integral for each subintervals** (with the trapezoidal method), and **adding the results**. (subintervals have identical width h)

$$I(f) = \int_a^b f(x)dx \approx \frac{1}{2} \sum_{i=1}^N [f(x_i) + f(x_{i+1})] (x_{i+1} - x_i)$$



$$I(f) \approx \frac{h}{2} \sum_{i=1}^N [f(x_{i+1}) + f(x_i)]$$

$$I(f) \approx \frac{h}{2} [f(a) + 2f(x_2) + 2f(x_3) + \dots + 2f(x_N) + f(b)]$$

$$I(f) \approx \frac{h}{2} [f(a) + f(b)] + h \sum_{i=2}^N f(x_i)$$

Example

Question: Consider $f(x) = 2 + \sin(2\sqrt{x})$. Use the composite trapezoidal rule with 11 sample points to compute an approximation to the integral of $f(x)$ taken over $[1, 6]$.

Simpson's methods

- The trapezoidal method described in the last section relies on approximating the integrand by a straight line. A better approximation can possibly be obtained by approximating the integrand with a nonlinear function that can be easily integrated.
- One class of such methods, called Simpson's rules or Simpson's methods, uses
 - quadratic (Simpson's 1/3 method), and
 - cubic (Simpson's 3/8 method)polynomials to approximate the integrand.

Simpson's 1/3 Method

- In this method, a quadratic (second-order) polynomial is used to approximate the integrand.
- The coefficients of a quadratic polynomial can be determined from **three points**.
- For an integral over the domain $[a, b]$, the three points used are the two endpoints $x_1 = a$, $x_3 = b$, and the midpoint $x_2 = (a + b)/2$.
- The polynomial can be written in the form:

$$p(x) = \alpha + \beta(x - x_1) + \gamma(x - x_1)(x - x_2)$$

where α , β , and γ are unknown constants evaluated from the condition that the polynomial passes through the points, $p(x_1) = f(x_1)$, $p(x_2) = f(x_2)$, and $p(x_3) = f(x_3)$.

Simpson's 1/3 Method

- These conditions yields

$$\alpha = f(x_1)$$

$$\beta = [f(x_2) - f(x_1)] / (x_2 - x_1)$$


$$\gamma = \frac{f(x_3) - 2f(x_2) + f(x_1)}{2(h)^2}$$

where $h = (b - a)/2$

- Substituting the constants back and integrating $p(x)$ over the interval $[a, b]$ gives

$$\begin{aligned} I = \int_{x_1}^{x_3} f(x)dx &\approx \int_{x_1}^{x_3} p(x)dx = \frac{h}{3} [f(x_1) + 4f(x_2) + f(x_3)] \\ &= \frac{h}{3} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \end{aligned}$$

Simpson's 1/3 Method




$$I = \frac{h}{3} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right]$$

- The name 1/3 in the method comes from the fact that there is a factor of 1/3 multiplying the expression in the brackets.
- As with the rectangular and trapezoidal methods, a more accurate evaluation of the integral can be done with a composite Simpson's 1/3 method.

Composite Simpson's 1/3 method

- The whole interval is divided into small subintervals. Simpson's 1/3 method is used to calculate the value of the integral in each subinterval, and the values are added together.






$$I(f) \approx \frac{h}{3} \left[f(a) + 4 \sum_{i=2,4,6}^N f(x_i) + 2 \sum_{j=3,5,7}^{N-1} f(x_j) + f(b) \right]$$

Composite Simpson's $1/3$ method

- It is important to point out that previous equation can be used only if two conditions are satisfied:
 - The subintervals must be equally spaced.
 - The number of subintervals within $[a, b]$ must be an even number

References

-  Numerical Methods Using MATLAB by Matthews and Fink, Pearson
-  Applied Numerical Methods with MATLAB for Engineers and Scientists, Third Edition
Steven C. Chapra, McGraw-Hill
-  Numerical Methods for Engineers and Scientists An Introduction with Applications using
MATLAB, Third Edition, Amos Gilat and Vish Subramaniam, John Wiley & Sons



Thank you!